

POLICY STATEMENT

Fluoridation of Public Drinking Water | Updated 21 August 2020



INTRODUCTION

Communities throughout the United States have used the fluoridation of drinking water supplies for over 70 years as a strategy to reduce tooth decay (dental caries)¹⁻⁷. The observed decline in national averages for the prevalence and severity of dental caries since the initiation of the program in 1945 has been deemed one of the greatest modern public health successes of the 20th century^{3, 5-9}. Despite these gains, a debate exists concerning the potential health risks of water fluoridation versus the observed benefit of the intervention^{1, 10, 11}. As a result, Public Health Madison & Dane County occasionally receives phone calls, emails, and letters from residents expressing concern about the fluoridation of local water supplies. Therefore, this document was produced and regularly reviewed to provide a brief overview of the current status of public concerns and an overview of the scientific literature.

OVERVIEW

Naturally occurring fluoride is found in all water supplies across the United States; the concentration is dependent upon the geology of the water body and the occurrence of fluoride-bearing minerals and materials^{1, 6, 8}. The discovery of the potential health applications of fluoride in the early 1930s led to the development of the first clinical trial of artificial fluoridation of community water supplies in Grand Rapids, MI in 1945. The trial was designed to last for 15 years prior to any potential recommendation for the expansion of water fluoridation to other communities; however, the popularity of the program led to its initiation in other cities the following year¹. The City of Madison, Wisconsin began the fluoridation of drinking water supplies in 1948.

EFFECTIVENESS OF WATER FLUORIDATION TO PREVENT DENTAL CARIES

Opponents of water fluoridation of community water supplies cite two major issues to question the effectiveness of the program; a comparable reduction in dental caries in non-fluoridated communities and the improved availability of fluoride-containing products make the treatment of community drinking water unnecessary.

Research has demonstrated that differences in the rate of dental caries in fluoridated and non-fluoridated communities have gradually decreased since the inception of water fluoridation programs. Opponents have used this and similar research to question the continued effectiveness of water fluoridation but this argument ignores the diffusion effect of fluoride containing products including food, beverages, dietary supplements, and dental products that were manufactured in fluoridated communities and sold in non-fluoridated areas^{1, 6, 8}.

Therefore, the non-fluoridated communities also experience an indirect benefit derived from

water fluoridation programs resulting in the comparable reduction of dental caries observed in these areas. This argument also ignores the larger benefit repeatedly reported among lower income and minority populations and the noted increase in dental caries among communities that have discontinued water fluoridation^{3, 5, 6, 7, 12 - 17}.

Oral hygiene has gradually improved in the United States over the past several decades, including the increased use of fluoride-containing dental products such as rinses, toothpaste, and topical gels¹²⁻¹⁶. However, disparities in use of and access to products and services promoting oral health remain, especially among low income and minority populations. The use of water fluoridation has provided an effective and cost efficient method to deliver preventative services to promote oral health to all residents within a community served by municipal water supplies, regardless of income, race or ethnicity, ability level, insurance status, or access to care^{3, 8, 12, 13, 17}. Currently, over 200 million Americans are served by drinking water supplies that contain the accepted levels of fluoride to reduce dental caries (approximately 75%); the Healthy People 2020 initiative calls for an expansion of this coverage to 80% of the population receiving drinking water from public water systems^{3, 6, 15, 16, 18}.

POTENTIAL HEALTH CONCERNS

The beneficial health effects of exposure to low concentrations of fluoride result from its ability to reduce tooth enamel solubility, decrease acid production of plaque-producing organisms, and promote the remineralization of the enamel^{1, 6, 16}. Efforts to prevent or remove water fluoridation in Dane County communities are most commonly motivated by concerns of fluorosis of teeth and bone, toxicity of fluoride, increased risk of hip fracture among the elderly, decreased IQ among children, hypothyroidism and cancer.

1. Acute fluoride toxicity

This condition has been reported when fluoridated drinking water supplies reach a level of 30ppm⁹. Due to the utilization of well-designed fail-safe equipment, proper maintenance and calibration, and appropriate operating procedures these overdosing incidents are rare in the United States. Symptoms normally occur within hours of exposure and include skin irritation, nausea, vomiting, diarrhea, and muscle weakness. Depending upon severity, observable symptoms resolve quickly following cessation of exposure. In severe cases, fluoride poisoning may result in cardiac arrest.¹⁹⁻²¹

2. Dental and skeletal fluorosis

These conditions are well-documented results of prolonged exposure to excess fluoride^{1, 9, 15, 16}. The development of dental fluorosis and skeletal fluorosis are attributed to the toxicokinetic properties of fluoride following exposure. Following ingestion, 75-90% of the compound is absorbed and readily distributed throughout the body⁹. Approximately 35-48% of the absorbed fluoride is retained by the body; an estimated 99% of the compound body burden is stored in the calcium rich areas of the bones and teeth (dentine and

enamel)^{9, 22}. This pattern of distribution and storage may lead to adverse impacts on the teeth and skeletal systems of individuals chronically exposed to excess natural and/or introduced levels of fluoride in public drinking water supplies^{1, 9, 15, 16, 22}.

Dental fluorosis is characterized by the staining and disruption of normal enamel formation of the teeth; the markings can range from unnoticeable in very mild cases (most common) to brown stains and pitting of the enamel in severe cases (rare)^{1, 9, 16, 23}.

Although severe cases can lead to brittle teeth and more teeth wear, all forms of dental fluorosis are considered by the United States Centers for Disease Control and Prevention (CDC) and the Department of Health and Human Services (DHHS) to be a cosmetic concern rather than an adverse health effect²³. Reported cases of dental fluorosis have increased in the United States since the widespread initiation of water fluoridation. Prior to the adoption of the program, the prevalence of dental fluorosis was 12-15%; modern rates of this condition have shown reported increases²⁴⁻²⁶. A study conducted by the CDC reported that an estimated 23% of persons aged 6 to 39 years had a very mild or greater dental fluorosis while approximately 32% of children and adolescents aged 6 to 19 years were reported with the condition. The risk of dental fluorosis development is limited to children 8 years of age or younger; tooth development occurs during this age range and the enamel is susceptible to the effects of fluoride when chronically exposed to levels of fluoride larger than 2 ppm. The current optimal level of fluoride recommended for drinking water sources is 0.7 ppm. Children older than 8 years, adolescents, and adults are not susceptible to dental fluorosis³.

Skeletal fluorosis is a disease characterized by increased density and brittleness of the skeletal system; the disease occurs in a range of severity dependent upon the level and duration of fluoride exposure. The mildest form of the disease can lead to arthritis-like symptoms including painful joints, limitations in movement, and reduced flexibility. Continual exposure to fluoride concentrations of 5ppm or greater may lead to osteosclerosis^{1, 9}. In the most severe cases, skeletal fluorosis can be a crippling disease, confining a patient to a wheelchair. The condition is extremely rare in the United States with only 5 confirmed cases reported in the last 35 years; each of these cases occurred in areas where natural fluoride levels were greater than 20ppm¹.

3. Hip fracture

Both excessive and inadequate intake of fluoride has been associated with an increased risk of hip fracture among the elderly^{1, 9-10}. However, research does not support an association between water fluoridation and increased rates of hip fracture^{16, 27, 28}. In fact, two systematic reviews of the literature published in 2000 and 2017 evaluating hip fracture and other types of bone fracture, respectively, also concluded that risk was not associated with water fluoridation¹⁶.

4. Reduction of IQ in children

Concerns have also been raised that water fluoridation is associated with lowered IQ and

other neurological effects from exposure. However, this concern is not supported by the science-based evidence. A number of systematic reviews of the literature and individual investigations does not support this association; including the recent response by the US Environmental Protection Agency (US EPA) to a TSCA Section 21 petition and a study released in 2018 conducted by the National Toxicology Program (NTP)^{15, 16, 29 - 31}. Despite this fact, some published investigations continue to suggest or imply neurological impacts among communities with fluoridated water at recommended levels^{15, 16, 29, 31 - 36}.

The human epidemiology studies that are often cited to support this concern are typically conducted in areas of the world with high naturally occurring levels of fluoride in the drinking water (e.g. China and Iran) and imply or are used by opponents of fluoridation to imply an association with community water fluoridation programs at recommended fluoride levels. In addition, limitations in the methodology of many of these studies lead to unsupported conclusions due to the impact of potential confounding factors that were not controlled or considered during the investigation^{29, 34, 35, 37, 38}.

One important limitation is that many of these studies use a cross-sectional design; a type of study design that does not establish causality because it cannot be determined if the health conditions occurred before or after the exposure in question. Cross-sectional study designs are best suited to help generate potential causal hypotheses further evaluated by more robust epidemiological investigations²⁹. Another important limitation of these studies is that potential founding factors are not adequately recognized and/or controlled which results in the inability to attribute the reported conclusions to fluoride exposure or other factors or exposures potentially present and not accounted for by the investigation. Examples include, but are not limited to, estimated water intake and not actual consumption, a lack of standardized testing of mental IQ, the failure to address other potential exposures that are associated with neurological impacts (e.g. lead and arsenic), exposure to other sources of fluoride (e.g. burning coal for heating), social inequities of the study population (e.g. poverty, nutritional deficiencies, and parental educational attainment), and/or measuring fluoride delivery in salt and imply association with water fluoridation^{15, 16, 28, 29, 31-36, 38-40}.

5. Hypothyroidism

An association has also been suggested between exposure to community water fluoridation and the incidence of hypothyroidism; a condition in which the thyroid gland doesn't produce a sufficient amount of hormones to function at a normal level⁴¹⁻⁴³. This suggested association is inconsistent with the literature and not supported by subsequent investigation of this potential association^{42, 44 - 46}.

Hypothyroidism is largely classified as an autoimmune disease (e.g. Hashimoto's thyroiditis) that is strongly associated with age (over 50 years) and sex (female). However, the condition can also be attributed to other underlying variables including the use of certain medications (e.g. lithium), both an increased or decreased intake of dietary iodine,

socioeconomic status, pregnancy, family history of thyroid disease, surgery, and radiation therapy^{16, 44, 46 - 48}. Many of the studies that report an association between water fluoridation at the recommended optimal level and hypothyroidism fail to account for many of these potential confounding factors that could result in spurious associations that are not reliable, representative, or generalizable to communities with fluoridated water resources.

6. Cancer

According to the World Health Organization (WHO), a large number of studies exploring the issue demonstrate no consistent evidence of any association between the consumption of controlled fluoridated drinking water with an increased risk of cancer^{9, 16}.

The majority of the concern about a potential cancer risk associated with the exposure to fluoridated drinking water is the development of osteosarcoma; a rare type of bone cancer typically diagnosed in children and teens in the United States. Similar to other types of cancers, the body of evidence does not display a consistent association between the consumption of drinking water fluoridated at recommended levels and the risk of osteosarcoma^{11, 49 - 52}. For example, research performed by the Harvard School of Public Health in 2006 reported that water fluoridation was associated with a higher risk of osteosarcoma in males but not females. However, early results from the second half of this investigation did not match the initial findings and the researchers advised caution in interpreting the results. The second part of the Harvard study was published in 2011 and found no association between water fluoridation and osteosarcoma risk^{50, 52}.

More recent studies have compared rates of osteosarcoma in areas of higher versus lower levels of water fluoridation in the United States, Great Britain, and Ireland; these studies have also not reported an increased risk in areas of water fluoridation⁵⁰.

FLUORIDATION COMPOUND SOURCES AND POTENTIAL CONTAMINATES

There are three basic compounds that are utilized for water fluoridation; sodium fluoride, sodium fluorosilicate, and fluorosilicic acid. Each of these compounds is derived from phosphorite rock, a source that is primarily used in the production of phosphate fertilizer. Phosphorite contains a mixture of calcium phosphate, calcium carbonate (limestone), and apatite; the mineral apatite contains approximately 3 to 7% fluoride overall and is considered the primary source of the fluoride used in water treatment^{1, 16}. The association of water fluoridation additives and the production of phosphate fertilizer have led to safety concerns by opponents of the intervention¹. The majority of these concerns center on potential impurities entering the drinking water supply as a result of the water fluoridation; specifically lead, arsenic, and radionuclides^{1, 16, 53}.

Regulatory processes are in place to protect community water supplies that either restricts and/or prevents the introduction of impurities from the fluoridation of drinking water. The U.S.

Environmental Protection Agency (US EPA) is responsible for the regulation of drinking water and to assure its safety in accordance with the Safe Drinking Water Act (SDWA). The SDWA requires that all additives used in water treatment plants, including fluoride additives, must meet strict regulatory standards in regards to their production, maintenance, and application. Each additive is subject to a system of standards, testing, and certification by the American Water Works Association (AWWA) and the National Sanitation Foundation/ American Standards Institute (NSF/ ANSI). Testing by the NSF for water quality has demonstrated that the vast majority of fluoride additive samples do not have detectable levels of arsenic derived from the addition of these compounds; water samples that do test positive are much lower than the EPA allowable levels. Other impurities, including lead and radionucleotides, are typically reported at levels lower than the detected arsenic levels^{22, 53, 54}. Aside from the testing of impurities, the recommended optimum fluoride concentration is 0.7 ppm; these levels are monitored to ensure appropriate concentrations are maintained in communities that fluoridate drinking water supplies^{3, 15, 16}.

The water fluoridation program for the City of Madison currently utilizes hydrofluorosilicic acid as its primary source for the fluoridation of community drinking water supplies. The compound is obtained from Hawkins Chemical, Inc. via an annual renewable contract. In addition to the federal requirements to ensure water quality, the City of Madison Water Utility has also designed and initiated additional safe guards to maintain safe water supplies. Standard operating procedures (SOPs) were designed in cooperation with Public Health Madison and Dane County to govern the operation of water fluoridation, routine maintenance of all equipment associated with the fluoridation process, and the daily monitoring of the water fluoride levels to ensure optimal recommended levels of fluoridation. Impurities, including potential impurities introduced by water fluoridation are also monitored in order to ensure that water quality standards are in accordance with regulatory policies; samples are derived from water entering the distribution center, which occurs after fluoridation to ensure the accurate reporting of water quality⁵⁵.

SUMMARY AND RECOMMENDATIONS

The occurrence of dental caries has been substantially reduced in the United States in recent decades, predominately through the widespread use of fluoride. Unfortunately, disparities among low income and minority populations are still quite prevalent^{2, 3, 9, 15, 16, 17}. This trend has also been reported in Wisconsin including Dane County; especially among populations of color. For example, between September 2014 and June of 2015, the Madison Metropolitan School District (MMSD) reported that 47% of students reporting urgent dental needs were African American children. In 2013, the rates for patients visiting hospital emergency departments in Dane County for dental pain were also higher for African Americans than for Whites, Hispanics, or other racial groups; similar findings were reported in 2015 providing further evidence of the persistence of oral health disparities^{56, 57}. These inequities underline the need for continued intervention efforts to address this inequity; water fluoridation is one of these tools.

To reach children and other at-risk populations for dental caries, water fluoridation is still the most efficient method of delivering safe and effective levels of fluoride. Therefore, Public

Health Madison and Dane County supports and recommends water fluoridation using the optimum fluoride concentration of 0.7 ppm as recommended by the United States Department of Health and Human Services^{3, 15, 16}. However, it should be recognized that drinking water fluoridation is a complex process that must be well monitored and controlled.

Careful review of the scientific literature and consultation with local and national experts has identified no evidence for adverse health impacts associated with water fluoridation at recommended levels. This fact and the continued high prevalence of dental caries and associated pain, expense and potentially serious medical consequences make the continuation and expansion of well controlled drinking water fluoridation a public health necessity in Dane County.

Prepared by: Jeffery S. Lafferty, Environmental Epidemiologist

Doug Voegeli, Director of Environmental Health

Janel Heinrich, Director of Public Health

REFERENCES

1. Ananian, A, Solomowitz, BH, and Dowrich, IA. (2006). Fluoride: a controversy revisited. *New York State Dental Journal*, 72(3), 14-18.
2. Centers for Disease Control and Prevention. (2007). CDC statement on the 2006 National Research Council (NRC) report on fluoride in drinking water. Retrieved on August 19, 2020 from: <http://kkw.org/wp-content/uploads/2016/06/Fluoride-CDC-reportpdf.pdf>
3. Centers for Disease Control and Prevention (2016). Water fluoridation basics. Retrieved on August 19, 2020 from: <https://www.cdc.gov/fluoridation/basics/index.htm>
4. Yeung, CA. (2007). Fluoride prevents caries among adults of all ages. *Evidence Based Dentistry*, 8(3), 72-73.
5. Griffen, S.O., Regnier, E., Griffen, P.M., and Huntley, V. (2007). Effectiveness of fluoride in preventing caries in adults. *Journal of Dental Research*, 86(5), 410 – 415.
6. American Dental Association. (2019). Fluoride in water. Retrieved on August 20, 2020 from: <https://www.ada.org/en/public-programs/advocating-for-the-public/fluoride-and-fluoridation>
7. Centers for Disease Control and Prevention. (2019). Over 70 years of community water fluoridation. Retrieved on August 21, 2020 from: <https://www.cdc.gov/fluoridation/basics/70-years.htm>
8. Centers for Disease Control and Prevention. (1999). Ten great public health achievements – United States, 1900-1999. *Morbidity and Mortality Weekly Report*, 48(12), 241-243.
9. Fawell, J, Bailey, K, Chilton, J, Dahi, E, Fewtrell, L, and Magara, Y. (2006). Fluoride in drinking water. The World Health Organization. IWA Publishing: London, UK.
10. Connett, P, Connett, E, and Connett, M. (2001). Fluoridation: time for a second look? Retrieved on August 21, 2020 from: <http://fluoridealert.org/news/fluoridation-time-for-a-second-look/>
11. Peckham, S. and Awofeso, N. (2014). Water fluoridation: A critical review of the physiological effects of ingested fluoride as a public health intervention.
12. Cho, H.J., Lee, H.S., Paik, D.I, and Bae, K.H. (2014). Association of dental caries with socioeconomic status in relation to different water fluoridation levels. *Community Dentistry and Oral Epidemiology*, 42(6), 536 – 542.
13. Burt, B.A. (2002). Fluoridation and social equity. *Journal of Public Health Dentistry*, 62(4), 195 – 200.
14. American Dental Association. (2015). Minority oral health in America: Despite progress, disparities persists. Retrieved on August 19, 2020 from: https://www.ada.org/~media/ADA/Advocacy/Files/160523_Kelly_Report_Dental_Chapter.pdf?la=en

15. Water Research Foundation (2015). State of science: Community water fluoridation. Retrieved on August 21, 2020 from: <https://www.waterrf.org/research/projects/state-science-community-water-fluoridation>
16. National Fluoridation Advisory Committee of the American Dental Association Council on Advocacy for Access and Prevention. (2018). *Fluoridation facts 2018*. American Dental Association.
17. McLaren, L, McNeil, DA, Potestio, M, Patterson, S, Thawer, S, Faris, P, Shi, C, and Shwart, L. (2016). Equity in children's dental caries before and after cessation of community water fluoridation: differential impact by dental insurance status and geographic material deprivation. *Internal Journal for Equity in Health*, 15:24, 1 – 9.
18. United States Department of Health and Human Services. (2019). Healthy People 2020: Topics and objectives: Oral Health. Retrieved on August 20, 2020 from: <https://www.healthypeople.gov/2020/topics-objectives/topic/oral-health/objectives#5003>
19. Hoffman, R, Mann, J, Calderone, J, Trumbull, J, and Burkhart, M. (1980). Acute fluoride poisoning in a New Mexico elementary school. *Pediatrics*, 65(5), 897-900.
20. Vogt, RL and Witherell, L. (1982). Acute fluoride poisoning associated with an on-site fluoridator in a Vermont elementary school. *American Journal of Public Health*, 72, 1168-1171.
21. Gessner, BD, Beller, M, Middaugh, JP, and Whitford, GM. (1994). Acute fluoride poisoning from a public water system. *New England Journal of Medicine*, 330(2), 95-99.
22. Casarett and Doull's Toxicology: The Basic Science of Poisons, Sixth Edition. Klaassen, C.D. (ed.). McGraw-Hill Publishing, New York, 2001.
23. United States Department of Health and Human Services. (2018). Statement on the evidence supporting the safety and effectiveness of community water fluoridation. Retrieved on August 21, 2020 from: <https://www.cdc.gov/fluoridation/pdf/Scientific-Statement-on-Community-Water-Fluoridation-h.pdf>
24. Clark, DC. (1994). Trends in prevalence of dental fluorosis in North America. *Community Dentistry and Oral Epidemiology*, 22(3), 148-152.
25. Heller, KE, Eklund, SA, and Burt, BA. (1997). Dental caries and dental fluorosis at varying water concentrations. *Journal of Public Health Dentistry*, 58(3), 136-143.
26. Beltran-Aguilar, E.D., Barker, L.K., Canto, M.T., Dye, B.A., Gooch, B.F., Griffin, S.O., ... and Wu, T.. (2005). Surveillance for dental caries, dental sealants, tooth retention, edentulism, and enamel fluorosis- United States, 1988 – 1994 and 1999-2002. *Morbidity and Mortality Weekly Report*, 54(03), 1-44.
27. Park, EY, Hwang, SS, Kim, JY, and Cho, SH. (2008). Effects of long-term fluoride in drinking water on risks of hip fracture of the elderly: an ecologic study based on database of hospitalization episodes. *Journal of Preventative Medicine and Public Health*, 41(3), 147-152.
28. Phipps, K. (1995). Fluoride and bone health. *Journal of Public Health Dentistry*, 55(1), 53-56.

29. Office of the Federal Register, National Archives and Records Administration. (2017). Fluoride chemicals in drinking water: TSCA Section 21 Petition: Reasons for agency response. *Federal Register*, 82(37), 11878 – 11890.
30. American Dental Association. (2018). Studies reaffirm safety of fluoridation. Retrieved on August 21, 2020 from: <https://www.ada.org/en/publications/ada-news/2018-archive/may/studies-reaffirm-safety-of-fluoridation>
31. McPherson, C.A., Zhange, G., Gilliam, R., Brar, S.S., Wilson, R., Brix, A., and Picut, C. (2018). An evaluation of neurotoxicity following fluoride exposure from gestational through adult ages in long-evans rats. *Neurotoxicity Research*, 34(4), 781 – 798.
32. Bashash, M., Thomas, D., Hu, H., Martinez-Mier, A., Sanchez, B.N., Basu, N., ... Hernandez-Avila, M. (2017). Prenatal fluoride exposure and cognitive outcomes in children at 4 and 6 – 12 years of age in Mexico. *Environmental Health Perspectives*, 125(9):097017. doi: 10.1289/EHP655
33. Choi, A. L., Sun, G., Zhang, Y., and Grandjean, P. (2012). Developmental fluoride neurotoxicity: A systematic review and meta-analysis. *Environmental Health Perspectives*, 120, 1362 – 1368.
34. Hong, F, Cao, Y, Yang, D, and Wang, H. (2001). Research on the effects of fluoride on child intellectual development under different environmental conditions. *Chinese Primary Health Care*, 15(3), 56 – 57. (translated report republished in *Fluoride*, 41(2), 156 – 160, April – June 2008).
35. Wang, SX, Wang, ZH, Cheng, XT, Li, J. Sang, ZP, Zhang, XD, ... and Wang, ZQ. (2007). Arsenic and fluoride exposure in drinking water: Children's IQ and growth in Shanyin County, Shanxi Province, China. *Environment Health Perspective*, 115(4), 643 – 647.
36. Green, R, Lanphear, B, Hornung, R, Flora, D, Martinez-Mier, A, Neufield, R, ... and Till, C. (2019). Association between maternal fluoride exposure during pregnancy and IQ scores in offspring in Canada. *Journal of the American Medical Association* 173(10), 940 – 948.
37. Dehghani, MA, Haghighat, GA, and Yousefi, M. (2018). Data on fluoride concentration in drinking water resources in Iran: A case study of Fars province; Larestan region. *Data Brief*, 19, 842 – 846.
38. Lu, Y, Sun, ZR, Wu, LN, Wang, X, Lu, W, Liu, SS.(2000). Effects on high fluoride water on intelligence in children. *Fluoride*, 33(2), 74 – 78.
39. Ando, M, Tadano, M, Yamamoto, S, Tamura, K, Asanuma, S, Watanabe, T. ... and Cao, S. (2001). Health effects of fluoride pollution caused by coal burning. *Science of the Total Environment*, 271(1-3), 107 – 116.
40. Li, F, Chen, X, Huang, R, and Xi, Y. (2009). The impact of endemic fluorosis caused by the burning of coal on the development of intelligence in children. *Journal of Environment and Health*, 26, 838 – 840.41. Peckham, S., Lowery, D., and Spencer, S. (2015). Are fluoride levels in drinking water associated with hypothyroidism prevalence in England? A large observational

study of GP practice data and fluoride levels in drinking water. *Journal of Epidemiology and Community Health*, 69, 619 -624.

42. American Thyroid Association. (2015). Is fluoridated water associated with a higher prevalence of hypothyroidism? Retrieved on August 21, 2020 from:

https://www.thyroid.org/wp-content/uploads/publications/ctfp/volume8/issue6/ct_public_v86_3.pdf

43. Kheradpisheh, Z., Mirzaei, M., Mahvi, A.H., Mokhtari, M., Azizi, R., Fallahzadeh, H., and Ehrampoush, M.H. (2018). Impact of drinking water fluoride on human thyroid hormones: A case control study. *Scientific Reports*, 8(2674), 1 – 7. doi: 10.1038/s41598-018-20696-4

44. Newton, J.N., Young, N., Verne, J., and Morris, J. 2015). Water fluoridation and hypothyroidism: results of this study need much more cautious interpretation. *Journal of Epidemiology and Community Health*, 69, 617 – 618.

45. Grimes, D.R. (2015). Commentary on: Are fluoride levels in drinking water associated with hypothyroidism prevalence in England? A large observational study of GP practice data and fluoride levels in drinking water. *Journal of Epidemiology and Community Health*, 69, 616.

46. Barberio, A.M., Hosein, F.S., Quinonez, C., and McLaren, L. (2017). Fluoride exposure and indicators of thyroid function in the Canadian population: implications for community water fluoridation. *Journal of Epidemiology and Community Health*, 71, 1019 – 1025.

47. American Thyroid Association. (2019). Hypothyroidism (underactive). Retrieved on August 20, 2020 from: <https://www.thyroid.org/hypothyroidism/>

48. World Health Organization. (2004). Iodine status worldwide: WHO global database on iodine deficiency. Retrieved on August 20, 2020 from: <https://www.who.int/vmnis/iodine/status/en/>

49. Comber, H., Deady, S., Montgomery, E., and Gavin, A. (2011). Drinking water fluoridation and osteosarcoma incidence on the island of Ireland. *Cancer Causes Control*, 22, 919 – 924.

50. American Cancer Society. (2019). Water fluoridation and cancer risk. Retrieved on August 21, 2020 from: <https://www.cancer.org/cancer/cancer-causes/water-fluoridation-and-cancer-risk.html>

51. Kim, F.M., Hayes, C., Williams, P.L., Whitford, G.M., Joshipura, K.J., Hoover, R.N., and Douglass, C.W. (2011). An assessment of bone fluoride and osteosarcoma. *Journal of Dental Research*, 90(10), 1171 – 1176.

52. National Cancer Institute. (2019). Fluoridated water. Retrieved on August 20, 2020 from: <https://www.cancer.gov/about-cancer/causes-prevention/risk/myths/fluoridated-water-fact-sheet>

53. Centers for Disease Control and Prevention. (2014). Water fluoridation additives. Retrieved on August 20, 2020 from: <https://www.cdc.gov/fluoridation/engineering/wfadditives.htm>

54. NSF International. (2019). Fact sheet on fluoridation products and fluoride. Retrieved on August 21, 2020 from: https://www.nsf.org/newsroom_pdf/Fluoride_Fact_Sheet_2019.pdf

55. Personal conversation with Joseph Grande, Water Quality Manager, City of Madison Water Utility.

56. Public Health Madison and Dane County. (2016). Oral health in Dane County. Retrieved on May 13, 2019 from: <http://www.publichealthmdc.com/documents/OralHealth201610.pdf>

57. Public Health Madison and Dane County. (2017). A report on Dane County hospital emergency department and urgent care visits for non-traumatic dental pain. Retrieved on May 13, 2019 from: <http://www.publichealthmdc.com/documents/ED-UCCDentalDataRpt201703.pdf>